



Fig. 1a.

Peptide pool 3



Gliadin digest



BEST AVAILABLE COPY

Fig.1b.




Fig.2a.





Fig.2b.



4/47




Fig.4a.





Fig.4b.



6/47

Fig.5.





Fig.6.



7/47

Fig.7a.

CD4 depletion






Fig.7b.


HLA-Class II restriction





9/47

Fig.9.



10/47

Fig.10.





Fig.11.




Fig.12a.

Dose response to A-gliadin 57-73 QE65:
QLQPFPPQPELPYPQPQS.




Fig.12b.

Dose response to GDA4_WHEAT P04724 84-100 QE92:
PQLPYPQPELPYPQPQP.




Fig.12c.

Dose response to A-gliadin 57-73:
QLQFPQPQLPYQPQPS (2.5, 25 & 250 mcg/ml),
and A-gliadin 57-73 (25 mcg/ml) + tTG treatment.




Fig.12d.

Dose response to GDA4_WHEAT P04724 84-100:
PQLPYQPQLPYQPQP (2.5, 25 & 250 mcg/ml),
and P04724 84-100 (25 mcg/ml) + tTG treatment.




Fig.12e.

Dose response to the DQ2-restricted α gliadin T cell epitope A-gliadin 57-68 QE65: QLQPFPPQPELPY (E65) (2.5, 25 & 250 mcg/ml), and A-gliadin 57-68: QLQPFPPQPQLPY (Q65) (25 mcg/ml) +/- tTG treatment.




Fig.12f.

Dose response to the DQ2-restricted α gliadin T cell epitope α -2 62-75 QE65 & QE72: PQPELPYPQPELPY (E65) (2.5, 25 & 250 mcg/ml), and α -2 62-75: PQPQLPYPQPQLPY (Q65) (25 mcg/ml) +/- tTG treatment.




Fig.12g.

Dose response to the DQ8-restricted α gliadin T cell epitope GDA9 202-219: QE208 & 216: QQYPSGEGSFQPSQENPQ (E) (25 & 250 mcg/ml), and to GDA9 202-219 QQYPSGQGSFQPSQQNPQ (Q) (25 mcg/ml) +/- tTG treatment.




Fig.12h.

Dose response to the DQ2-restricted γ gliadin T cell epitope GDB2 134-153 QE140, 148,150: QQLPQPEQPQQSFPEQERPF (E) (25 & 250 mcg/ml), and to GDB2 134-153: QQLPQPQQPQQSFPPQQRPF (Q) (25 mcg/ml) +/- tTG treatment.




Fig.13a.

Dose response to gliadin digest by
chymotrysin.




Fig.13b.

Dose response to gliadin digested by
chymotrysin then treated with tTG.





Fig. 13c.

Total ELISpot responses to A-gliadin 57-73 QE65 (25mcg/ml) versus A-gliadin 57-73 QE65 responses as percent of tTG gliadin (500mcg/ml) responses.


Responses to dominant epitope and complete antigen (A-gliadin 57-73 QE65 and tTG-treated gliadin)



(Fig.14.)


Bioactivity of gliadin polymorphisms of A-gliadin 57-73
(A) in coeliac subjects 6/7 days after gluten challenge
(Gamma-Interferon Elispot) (n=4).

Fig.14a. Unmodified (25 mcg/ml): Means (error bars: 1 SEM)



A	QLQPFPPQPQLPYQPQS	I	QLQPFPPQPQLSYSQPQP
B	QLQPFPPQPQLPYQPQP	J	QPQPFPPPPQLPYPQTQP
C	QLQPFPPQPQLPYQPQL	K	PQLPYQPQLPYQPQP
D	QLQPFPPQPQLPYLQPQS	L	PQLPYQPQLPYQPQL
E	QLQPFPPRPQLPYQPQP	M	PQPQFPPQLPYQPQS
F	QLQPFPPQPQLPYSQPQP	N	PQPQFPPQLPYQPQS
G	QLQPFLLQPQLPYSQPQP	O	PQPQFPPQLPYPQTQP
H	QLQPFSSQPQLPYSQPQP	P	PQPQFPPQLPYQPQP

Fig.14b.




A QLQPFPPQPQLPYQPQS
 B QLQPFPPQPQLPYQPQP
 C QLQPFPPQPQLPYQPQL
 D QLQPFPPQPQLPYLQPQS
 E QLQPFPPQPQLPYQPQP
 F QLQPFPPQPQLPYSQPQP
 G QLQPFLLQPQLPYSQPQP
 H QLQPFSSQPQLPYSQPQP

I QLQPFPPQPQLSYSQPQP
 J QPQPFPPPPQLPYPQTQP
 K PQLPYQPQPQLPYQPQP
 L PQLPYQPQPQLPYQPQL
 M PQPQPFLPQLPYQPQS
 N PQPQPFPPQLPYQPQS
 O PQPQPFPPQLPYPQTQP
 P PQPQPFPPQLPYQPQP

Fig.14c.

QE65 substituted (25 mcg/ml): Means (error bars: 1 SEM)



A QLQPFPPQPQLPYQPQS
 B QLQPFPPQPQLPYQPQP
 C QLQPFPPQPQLPYQPQL
 D QLQPFPPQPQLPYLQPQS
 E QLQPFPPQPQLPYQPQP
 F QLQPFPPQPQLPYSQPQP
 G QLQPFLLQPQLPYSQPQP
 H QLQPFLLQPQLPYSQPQP

I QLQPFPPQPQLSYSQPQP
 J QPQPFFPPQLPYPQTQP
 K PQLPYQPQLPYQPQP
 L PQLPYQPQLPYQPQL
 M PQPQFFLPQLPYQPQS
 N PQPQFFPPQLPYQPQS
 O PQPQFFPPQLPYPQTQP
 P PQPQFFPPQLPYQPQP

Fig.14d. QE65-substituted (2.5 mcg/ml): Means (error bars: 1 SEM)





Fig.14e. QE65-substituted (250 mcg/ml): Means (error bars: 1 SEM)



A QLQPFPPQQLPYPQPQS
 B QLQPFPPQQLPYPQPQP
 C QLQPFPPQQLPYPQPQL
 D QLQPFPPQQLPYLQPQS
 E QLQPFPPQQLPYPQPQP
 F QLQPFPPQQLPYSQPQP
 G QLQPFLLQQLPYSQPQP
 H QLQPFSSQPQLPYSQPQP

I QLQPFPPQQLSYSQPQP
 J QPQPFPPQQLPYPQITQP
 K PQLPYPQPQLPYPQPQP
 L PQLPYPQPQLPYPQPQL
 M PQQQPFLLPQLPYPQPQS
 N PQQQPFPPQQLPYPQPQS
 O PQQQPFPPQQLPYPQITQP
 P PQQQPFPPQQLPYPQPQP

Fig.15.

Alanine scan: Means (error bars: 95% CI for mean)




Fig.16.

Lysine scan: Means (error bars: 95% CI for mean)




Fig.17.

Agonist activity of A-gliadin 57-73 QE65 variants according to position substituted (Mean of 8 coeliac subjects' PBMC responses in interferon gamma ELISPOT after gluten challenge)

QLQPF⁶⁰FPQPELPYPQPQS

60.....70





Fig.18.

Agonist activity of A-gliadin 57-73 QE65 variants according to position substituted (Mean of 8 coeliac subjects' PBMC responses in interferon gamma ELISPOT after gluten challenge)

QLQPFPPQPELPYPQPQS

60.....70



27/47

Fig.19.

Agonist activity of A-gliadin 57-73 QE65 variants according to position substituted (Mean of 8 coeliac subjects' PBMC responses in interferon gamma ELISPOT after gluten challenge)

QLQPFPPQPELPYPQPQS

60.....70




Fig.20.

Agonist activity of A-gliadin 57-73 QE65 variants according to position substituted (Mean of 8 coeliac subjects' PBMC responses in interferon gamma ELISPOT after gluten challenge)

QLQPF⁶⁰QPELPYPQPQS⁷⁰

60.....70




Fig.21.

Agonist activity of A-gliadin 57-73 QE65 variants according to position substituted (Mean of 8 coeliac subjects' PBMC responses in interferon gamma ELISPOT after gluten challenge)

QLQPFPPQPELPYPQPQS

60.....70




Fig.22.

Agonist activity of A-gliadin 57-73 QE65 variants according to position substituted (Mean of 8 coeliac subjects' PBMC responses in interferon gamma ELISPOT after gluten challenge)

QLQPFPPQPELPYPQPQS
60.....70




Fig.23.

Agonist activity of A-gliadin 57-73 QE65 variants according to position substituted (Mean of 8 coeliac subjects' PBMC responses in interferon gamma ELISPOT after gluten challenge)

QLQPF⁶⁰PQPELPYPQPQS⁷⁰

60.....70




Fig.24.

Agonist activity of A-gliadin 57-73 QE65 variants according to position substituted (Mean of 8 coeliac subjects' PBMC responses in interferon gamma ELISPOT after gluten challenge)

QLQPFPPQPELPYPQPQS

60.....70




Fig.25.

Agonist activity of A-gliadin 57-73 QE65 variants according to position substituted (Mean of 8 coeliac subjects' PBMC responses in interferon gamma ELISPOT after gluten challenge)

QLQPFPPQPELPYPQPQS

60.....70




Fig.26.

Agonist activity of A-gliadin 57-73 QE65 variants according to position substituted (Mean of 8 coeliac subjects' PBMC responses in interferon gamma ELISPOT after gluten challenge)

QLQPFPPQPELPYPQPQS

60.....70





Fig.27.

Agonist activity of A-gliadin 57-73 QE65 variants according to position substituted (Mean of 8 coeliac subjects' PBMC responses in interferon gamma ELISPOT after gluten challenge)

QLQPF⁶⁰PQPELPYPQPQS

60.....70



(Fig.28.)

Interferon gamma ELISpot responses in newly diagnosed and treated coeliac subjects, before and after gluten challenge.

Fig.28a. Untreated, newly diagnosed coeliacs (Mean+SEM, n=9)




Fig.28b.




Fig.28c.




Fig.28d.




Fig.28e.




Fig.28f.




Fig.28g.






Figure 29. Bioactivity of prolamin homologues of A-gliadin 57-73 (IFNg-ELISpot, mean+SEM, n=6)



Omega: AAG17702 (141-157), Gamma: P21292 (96-112), SEC1: Q43639 (335-351), C-HOR: Q40055 (166-182). E-substituted peptides were synthesized with E for Q at position 9.

10/516837

Figure 30. Healthy HLA-DQ2 Subjects: Change in IFNgamma ELISpot
Responses to tTG-deamidated Gliadin Peptide Pools
(median change Day 6 vs Day 0, n=10)



42 / 47

Figure 31. Coeliac HLA-DQ2 Subjects: Change in IFNgamma ELISpot
Responses to tTG-deamidated Gliadin Peptide Pools
(median change Day 6 vs Day 0, n=6)




Figure 32. Individual Peptide Contributions to "Summed"
Gliadin Peptide Response





Figure 33. Coeliac HLA-DQ2/8 Subject C08: Gluten challenge induced IFNgamma ELISpot Responses to tTG-deamidated Gliadin Peptide Pools



45/47

Figure 34. Coeliac HLA-DQ2/8 Subject C07: Change in IFNgamma ELISpot Responses to tTG-deamidated Gliadin Peptide Pools






Figure 35. Coeliac HLA-DQ8/7 Subject C12: Gluten challenge induced IFNgamma ELISpot Responses to tTG-deamidated Gliadin Peptide Pools



47 / 47

Figure 36. Coeliac HLA-DQ6/8 Subject C11: Change in IFNgamma ELISpot Responses to tTG-deamidated Gliadin Peptide Pools



**This Page is Inserted by IFW Indexing and Scanning
Operations and is not part of the Official Record**

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

☐ **BLACK BORDERS**

☐ **IMAGE CUT OFF AT TOP, BOTTOM OR SIDES**

☐ **FADED TEXT OR DRAWING**

☒ **BLURRED OR ILLEGIBLE TEXT OR DRAWING**

☐ **SKEWED/SLANTED IMAGES**

☐ **COLOR OR BLACK AND WHITE PHOTOGRAPHS**

☐ **GRAY SCALE DOCUMENTS**

☐ **LINES OR MARKS ON ORIGINAL DOCUMENT**

☐ **REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY**

☐ **OTHER:** _____

IMAGES ARE BEST AVAILABLE COPY.

As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.